

Anisotropic Brane Cosmology with Variable G and Λ

Soma Nath, Subenoy Chakraborty,^{*} and Ujjal Debnath[†]

Department of Mathematics, Jadavpur University, Calcutta-32, India.

(Dated: February 7, 2008)

In this work, the cosmological implications of brane world scenario are investigated when the gravitational coupling G and the cosmological term Λ are not constant but rather there are time variation of them. From observational point of view, these time variations are taken in the form $\frac{\dot{G}}{G} \sim H$ and $\Lambda \sim H^2$. The behavior of scale factors and different kinematical parameters are investigated for different possible scenarios where the bulk cosmological constant Λ_5 can be zero, positive or negative.

PACS numbers: 04.50 +h , 04.20.Cv

I. INTRODUCTION

There are two fundamental physical parameters namely the gravitational coupling G and the cosmological term Λ , in the Einstein's theory of general relativity. Usually these parameters are assumed to be constants. But recent experimental and (or) observational evidences suggest that it is not unlikely to have small variations [1] of these fundamental parameters. However, alternative ideas about the variability of these parameters have been started long ago. The idea of variable G was first introduced by Dirac [2] in cosmological consideration and then subsequently by Sciama [3], Jordan [4] and others. Later, Brans and Dicke [5] proposed an extension (or modification) of Einstein's theory of gravity by introducing a scalar field ϕ . In this theory, the Newton's gravitational constant G is a variable and is related to the scalar field $\phi \sim G^{-1}$.

The cosmological term Λ on the otherhand, was originally suggested by Einstein himself to obtain static solution of his field equations. But after his realization that the universe is expanding, he discarded the cosmological term. Afterwards, the cosmological term has been introduced and subsequently rejected several times for various reasons. But recent measurements of the CMB anisotropy and the observations from type Ia supernovae demand a significant and positive cosmological constant [6, 7]. Also observations of gravitational lensing indicate the presence of a non-zero Λ . However, to resolve the cosmological constant problem (i.e., its observed value is about 120 orders of magnitude below the value for the vacuum energy density predicted by quantum field theory) a phenomenological solution namely a dynamical $\Lambda(t)$ has been suggested by several authors [8], arguing that Λ relaxes its present estimate due to the expansion of the universe. Also due to these recent observational predictions the vacuum energy density ρ_Λ ($= \Lambda/8\pi G$) has played an important role and the cosmological constant problem has been shifted to the *coincidence problem* namely why ρ and ρ_Λ happen to be of the same order of magnitude precisely at this very moment. To resolve this naturalness several quintessence models [9] have been proposed where the cosmological constant becomes a time dependent quantity. Recently Bonanno et al [10] (see also Shapiro et al [11]) have formulated a general frame work for cosmologies in which both Newton's constant and the cosmological constant are time dependent and have shown that ρ and ρ_Λ are approximately equal in the late universe. They have assumed that there exists an infrared attractive fixed point for the renormalization group flow of the (dimensionless) Newton's constant and cosmological constant respectively. Vishwakarma [12] has proposed a variable Λ proportional to H^2 (H = Hubble parameter) and has showed a good agreement with recent supernovae observations. Also this choice provides expected

^{*}Electronic address: subenoyc@yahoo.co.in

[†]Electronic address: ujjaldebnath@yahoo.com

large age of the universe.

In this paper, we analyse a 5D brane world model with a time dependent brane tension λ and constant 5D cosmological constant Λ_5 . We shall show that this will lead to a 4D effective theory with time dependent G and Λ . Although the fundamental theory is not yet known, the time dependent of the self-energy of the brane λ can be the result of some higher dimensional effect, such as a particular bulk-brane interaction, or the response of the brane to the change in the inter-brane distance in a 2-brane world model. Thus here we attempt to make a “phenomenological” study of the implications of such a time-dependence of λ on observable 4-dimensional parameters, such as G and Λ , and its consistency with current observational constraints.

II. BASIC EQUATIONS IN BRANE COSMOLOGY

The idea of brane world scenario was proposed by Randall and Sandrum [13]. They have shown that in a five dimensional space-time (called *bulk*) it is possible to confine the matter fields in a four dimensional hypersurface (called *3-brane*). The effective action in the bulk is [14]

$$A = \int d^5x \sqrt{-g_5} \left(\frac{1}{2\kappa_5^2} R_5 - \Lambda_5 \right) + \int_{\chi=0} d^4x \sqrt{-g} \left(\frac{1}{2\kappa_5^2} K^\pm + \lambda + \mathcal{L}_{matter} \right) \quad (1)$$

Here any quantity in the bulk is marked with subscript ‘5’ with $\kappa_5^2 (= \frac{8\pi G_5}{c^4})$ as the 5D gravitational coupling constant. The co-ordinates on the bulk are denoted by x^a , $a = 0, 1, 2, 3, 4$ while those on the brane as x^μ , $\mu = 0, 1, 2, 3$. So $\chi = x^4 = 0$ the four dimensional hypersurface, is defined as the brane world. The parameter λ stands for the brane tension which is assumed to be positive to recover the conventional gravity on the brane. The other parameter Λ_5 is the negative vacuum energy and is the only source of gravitational field on the bulk while $K^\pm$ are the intrinsic curvature on either side of the brane (characterized by $\chi > 0$ or < 0). Now the effective Einstein equations on the bulk are [15, 16]

$$G_{ab}^5 = -\Lambda_5 g_{ab}^5 + \kappa_5^2 T_{ab}^5(\text{brane}) \quad (2)$$

where $T_{ab}^5(\text{brane})$ (with $T_{ab}^5(\text{brane}) \cdot n^a = 0$) is the total energy-momentum tensor (vacuum+matter) on the brane i.e.,

$$T_{ab}^5(\text{brane}) = \delta(\chi) [\lambda g_{ab} + T_{ab}^{(m)}] = \delta(\chi) \delta_a^\mu \delta_b^\nu \tau_{\mu\nu} \quad (3)$$

Here $T_{ab}^{(m)}$ is usual matter energy tensor and the Dirac delta function indicates the fact that matter is confined in the space-like hypersurface $\chi = 0$ (the 3-brane) with induced metric g_{ab} .

From the Israel’s junction conditions

$$K_{\mu\nu}|_{\chi>0} - K_{\mu\nu}|_{\chi<0} = \kappa_5^2 \left(\tau_{\mu\nu} - \frac{1}{3} g_{\mu\nu} \tau \right)$$

Due to Z_2 -symmetry of the bulk,

$$K_{\mu\nu}|_{\chi>0} = -K_{\mu\nu}|_{\chi<0} = \frac{1}{2} \kappa_5^2 \left(\tau_{\mu\nu} - \frac{1}{3} g_{\mu\nu} \tau \right) \quad (4)$$

Hence we obtain

$$\tau_{\mu\nu} = \frac{2}{\kappa_5^2} (K_{\mu\nu} - g_{\mu\nu} K)$$

So from (4) we have

$$K_{\mu\nu}|_{\chi>0} = -\frac{1}{2}\kappa_5^2 \left[T_{\mu\nu}^{(m)} - \frac{1}{3}g_{\mu\nu}(T^{(m)} + \lambda) \right]$$

Then the effective Einstein equations on the brane are ([15]; see also [17]) (choosing $8\pi = 1, c = 1$)

$$G_{\mu\nu} = -\Lambda g_{\mu\nu} + \kappa^2 T_{\mu\nu}^{(m)} + \kappa_5^4 S_{\mu\nu} - E_{\mu\nu} \quad (5)$$

(Note that Greek indices refer to brane world while English small letters to bulk quantities).

The above Einstein equation has two correction terms to the energy-momentum tensor namely the local correction term $S_{\mu\nu}$ with expression

$$S_{\mu\nu} = \frac{1}{12}T^{(m)}T_{\mu\nu}^{(m)} - \frac{1}{4}T_{\mu}^{\alpha(m)}T_{\nu\alpha}^{(m)} + \frac{1}{24}g_{\mu\nu}(3T^{\alpha\beta(m)}T_{\alpha\beta}^{(m)} - T^{2(m)}) \quad (6)$$

while the other correction term is the non-local effects from the free gravitational field in the bulk with expression

$$E_{ab} = C_{abcd}n^cn^d \quad (7)$$

Here C_{abcd} is the usual Weyl tensor in the bulk, n^a the unit normal to the hypersurface $\chi = 0$ and the four dimensional cosmological constant Λ has the expression

$$\Lambda = \frac{\kappa_5^2}{2}(\Lambda_5 + \frac{\kappa_5^2\lambda^2}{6}) \quad (8)$$

with

$$\kappa^2 = \frac{\kappa_5^4\lambda}{6} \quad (9)$$

as the four dimensional gravitational constant. Now using the conservation of the total matter energy-momentum tensor on the brane namely

$$\tau_{\mu;\nu}^{\nu} = 0 \quad (10)$$

into the modified Einstein equations on the brane we have a constraint on $S_{\mu\nu}$ and $E_{\mu\nu}$ as

$$(E_{\mu}^{\nu} - \kappa_5^4 S_{\mu}^{\nu});_{\nu} = 0 \quad (11)$$

Maartens has shown that $E_{\mu\nu}$ can be decomposed as [16, 17]

$$E_{\mu\nu} = -\frac{6}{\kappa^2\lambda}[U(u_{\mu}u_{\nu} + \frac{1}{3}h_{\mu\nu}) + 2u_{(\mu}Q_{\nu)} + P_{\mu\nu}] \quad (12)$$

where the prefactor $\frac{6}{\kappa^2\lambda}$ (i.e., $\frac{\kappa_5^4}{\kappa^4}$) is introduced for dimensional reasons (note that $\lambda^{-1} \rightarrow 0$ gives the general relativistic limit) and the energy flux Q_{μ} and energy stresses $P_{\mu\nu}$ has the following properties :

(i) Q_{μ} is a spatial vector (i.e., $Q_{\mu}u^{\mu} = 0$)

(ii) $P_{\mu\nu}$ is a spatial (i.e., $P_{\mu\nu}u^{\nu} = 0$), symmetric ($P_{(\mu\nu)} = P_{\mu\nu}$) and trace-free ($P_{\mu}^{\mu} = 0$) tensor.

The scalar term U is called dark energy density as it has the same form as the energy momentum tensor of a radiation perfect fluid. Also u^μ is the 4-velocity on the brane with $h_{\mu\nu}$, the projection tensor orthogonal to u^μ on the brane. Further, using the constraint equation (11) between $E_{\mu\nu}$ and $S_{\mu\nu}$, it is possible to have evolution equations for U and Q_μ but not for $P_{\mu\nu}$. So the system of equations on the brane is not, in general closed.

As we are considering Bianchi models which have a simple transitive three dimensional group of isometries G_3 on the space-like hypersurfaces, so the cosmological models are spatially homogeneous with proper time as the only dynamical variables. Thus the metric on the brane can be written as [18]

$$ds^2 = -dt^2 + \alpha_i^2(t) dx^i dx^i \quad (13)$$

Further, the physical parameters which are of observational interest in cosmology are the following:

Expansion scalar: $\theta = 3H = \sum_{i=1}^3 H_i = \sum_{i=1}^3 \frac{\dot{\alpha}_i}{\alpha_i}$

Shear scalar:

$$\sigma^2 = \frac{1}{2} \sigma_{\mu\nu} \sigma^{\mu\nu} = \frac{3}{2} A H^2 \quad (14)$$

Deceleration parameter: $q = -\frac{(\dot{H} + H^2)}{H^2}$

where the mean anisotropy parameter A has the expression

$$A = \frac{1}{3} \sum_{i=1}^3 \left(1 - \frac{H_i}{H} \right)^2$$

and H_i and H are the directional Hubble parameters and mean Hubble parameters respectively.

Lastly, for Bianchi I and V model we get $Q_\mu = 0$ but $P_{\mu\nu}$ remains unrestricted. As there is no way of fixing the dynamics of this tensor, we shall study the particular case $P_{\mu\nu} = 0$. Then from the constraint equation (11) the dark energy density U is a function of the proper time alone i.e., $U = U(t)$ and the evolution equation for U takes the form

$$\dot{U} = -4HU \quad (15)$$

which on integration gives

$$U = U_0 V^{-\frac{4}{3}} \quad (16)$$

where $V = \prod_{i=1}^3 \alpha_i$ is the volume scale factor and U_0 is an integration constant.

III. BIANCHI I MODEL

The line element of a Bianchi I space-times generalizes flat FRLW metric to the anisotropic case as

$$ds^2 = -dt^2 + a_1^2(t) dx^2 + a_2^2(t) dy^2 + a_3^2(t) dz^2 \quad (17)$$

We have taken perfect fluid as the matter on the brane with expression

$$T_{\mu\nu}^{(m)} = (\rho + p) u_\mu u_\nu + p g_{\mu\nu} \quad (18)$$

where the energy density ρ and thermodynamic pressure p satisfy the isothermal equation of state namely

$$p = (\gamma - 1)\rho, \quad 1 \leq \gamma \leq 2. \quad (19)$$

The combinations of the non-vanishing components of the Einstein field equations on the brane (i.e.(5)) are [18]

$$3\dot{H} + \sum_{i=1}^3 H_i^2 = \Lambda - \frac{(3\gamma - 2)}{2} \kappa^2 \rho - \frac{(3\gamma - 1)}{12} \kappa_5^4 \rho^2 + \frac{U_0}{V^{\frac{4}{3}}} \quad (20)$$

and

$$\frac{1}{V} \frac{d}{dt} (V H_i) = \Lambda - \frac{(\gamma - 2)}{2} \kappa^2 \rho - \frac{(\gamma - 1)}{12} \kappa_5^4 \rho^2 + \frac{U_0}{V^{\frac{4}{3}}} \quad (21)$$

It is to be noted that so far we have not considered any variation of the 4D physical parameters namely G and Λ . The variation of G and Λ shows a time variation of brane tension λ (see equations (8) and (9)), keeping 5D quantities as invariant. From observational point of view it is generally assumed that the time variation of G can be expressed in terms of the Hubble parameter H as [19, 20]

$$\frac{\dot{G}}{G} = 3gH \quad \text{i.e., } G = G_0 V^g$$

where the dimensionless parameter g has present observational bound $|g| \leq 0.1$ and G_0 is constant of integration.

As from equation (9) $G \propto \lambda$ so the differential equation in λ has the same form namely,

$$\frac{\dot{\lambda}}{\lambda} = 3gH \quad \text{i.e., } \lambda = \lambda_0 V^g \quad (22)$$

where λ_0 is integration constant. Also the integration constants (G_0 and λ_0) are connected by the relation (using eq.(9)) $G_0 = \frac{\kappa_5^4}{6} \lambda_0$.

As a consequence, the energy conservation equation (10) becomes

$$\dot{\rho} + 3\gamma\rho H = -\lambda_0 g V^{g-1} \dot{V} \quad (23)$$

which on integration gives

$$\rho = D V^{-(1+\beta)} \quad (24)$$

with $\lambda_0 = F_0 D$ and $1 + \beta = -g = \frac{\gamma}{F_0 + 1}$.

Hence from observation β is restricted to $-1 < \beta \leq -0.966$. Also from equation (9) the brane tension is proportional to G and we write

$$\lambda = \frac{D(\gamma - 1 - \beta)}{(1 + \beta)} V^{-(1+\beta)} \quad (25)$$

Now using relations (8), (16), (22) and (23) in the field equation (21) we have

$$\frac{1}{V} \frac{d}{dt} (V H_i) = \frac{\kappa_5^2 \Lambda_5}{2} - \frac{\kappa_5^4 D^2 \gamma^2 \beta}{12(1 + \beta)^2} V^{-2(1+\beta)} + \frac{U_0}{V^{\frac{4}{3}}} \quad (26)$$

So adding equation (26) for $i = 1, 2, 3$ we get

$$\frac{1}{V} \frac{d}{dt} (V H) = \frac{1}{V} \frac{d}{dt} (V H_i) \quad (27)$$

which on integration gives

$$H = H_i + \frac{K_i}{V} \quad (28)$$

The integration constants K_i , $i = 1, 2, 3$ are restricted by the relation $\sum_{i=1}^3 K_i = 0$.

Combining equation (26) with (27) we have the differential equation in V (after integrating once) as (assuming $(1 + \beta) \neq 0$)

$$\dot{V}^2 = aV^2 + bV^{-2\beta} + cV^{\frac{2}{3}} + d_1 \quad (29)$$

or in integral form

$$t - t_0 = \int \frac{dV}{\sqrt{aV^2 + bV^{-2\beta} + cV^{\frac{2}{3}} + d_1}} \quad (30)$$

where $a = \frac{3}{2}\kappa_5^2\Lambda_5$, $b = \frac{\kappa_5^4 D^2 \gamma^2}{4(1+\beta)^2}$, $c = 9U_0$ and (d_1, t_0) are integration constants.

Also the scale factors can be obtained integrating once equation (28) and using equation (29) as

$$a_i = a_{i0} V^{\frac{1}{3}} \exp \left[\frac{K_i}{3} \int \frac{dz}{\sqrt{a + bz^{2(\beta+1)}}} \right], \quad i = 1, 2, 3$$

with a_{i0} 's as integration constants and $z = V^{-1}$. Further, if we assume the variation of the cosmological term (as mentioned earlier) as

$$\Lambda = \xi H^2 \quad (31)$$

(with ξ , a function of time) then using (25) and (31) in equation (8) we get a differential equation in V as

$$\dot{V}^2 = \frac{9}{2\xi}\kappa_5^2\Lambda_5 V^2 + \frac{3}{4\xi}\kappa_5^4 D^2 \frac{(\gamma - 1 - \beta)^2}{(1 + \beta)^2} V^{-2\beta}$$

Now comparing with equation (29) we have

$$a \left(1 - \frac{3}{\xi} \right) V^{2(1+\beta)} + \frac{b}{\gamma^2} \left[\gamma^2 - \frac{3}{\xi} (\gamma - 1 - \beta)^2 \right] + cV^{2(\frac{1}{3}+\beta)} + d_1 V^{2\beta} = 0 \quad (32)$$

which is true for all time and for all values of V . As powers of V in the 3rd and 4th term are in no way comparable to the first two terms for any choice of β so we can choose the integration constant $d_1 = 0$ and also $c = 0$. We note that the constant ' c ' is related to the non-local energy correction term (corresponds to an effective radiation) which is constrained to be small enough at the time of nucleosynthesis and it should be negligible today.

Moreover as a solution of (32) we have (in addition to $c = 0 = d_1$) three possibilities namely

$$(i) \quad \xi = 3 \quad \text{and} \quad \gamma^2 = (\gamma - 1 - \beta)^2 \quad \text{i.e.} \quad \gamma = \frac{(1 + \beta)}{2} \quad (33)$$

$$(ii) \quad a = 0 \quad \text{and} \quad \xi = \frac{3(\gamma - 1 - \beta)^2}{\gamma^2} \quad (34)$$

and

$$(iii) \quad \xi \text{ is a function of } V.$$

It is to be noted that the first choice contradicts the observational bounds for β [19] (i.e., $-1 < \beta \leq -0.966$), so only the second and the third possibilities will be discussed in the following for all possible values of Λ_5 namely, (i) $\Lambda_5 > 0$, (ii) $\Lambda_5 < 0$ and (iii) $\Lambda_5 = 0$.

Case I: Λ_5 (i.e., a) > 0

Here we have the solution of V from equation (30) as (the third possibility i.e., ξ is a function of V , is the only choice)

$$V^{(1+\beta)} = \sqrt{\frac{b}{a}} \sinh[(1+\beta)\sqrt{a}(t-t_0)] \quad (35)$$

which simplifies to (choosing $t_0 = 0$, so the big bang singularity occurs at $t = 0$)

$$V^{1+\beta} = \frac{D \kappa_5 \gamma}{(1+\beta)\sqrt{6\Lambda_5}} \sinh[(1+\beta)\sqrt{\frac{3\Lambda_5}{2}} \kappa_5 t] \quad (36)$$

with

$$\xi = \frac{3(\gamma - \beta - 1)^2}{\gamma^2} + \frac{\kappa_5^2 \Lambda_5}{2H^2} \left[1 - \frac{(\gamma - \beta - 1)^2}{\gamma^2} \right]$$

The expressions for different physical parameters are

$$\left. \begin{aligned} H &= \frac{\theta}{3} = \sqrt{\frac{\Lambda_5}{6}} \kappa_5 \coth[(1+\beta)\sqrt{\frac{3\Lambda_5}{2}} \kappa_5 t] \\ \rho &= \frac{\sqrt{6\Lambda_5}(1+\beta)}{\kappa_5 \gamma} \operatorname{cosech}[\sqrt{\frac{3\Lambda_5}{2}} \kappa_5 (1+\beta) t] \\ q &= 3(1+\beta) \operatorname{sech}^2[\sqrt{\frac{3\Lambda_5}{2}} \kappa_5 (1+\beta) t] - 1 \\ \lambda &= \frac{(\gamma - \beta - 1)\sqrt{6\Lambda_5}}{\kappa_5 \gamma} \operatorname{cosech}[\sqrt{\frac{3\Lambda_5}{2}} \kappa_5 (1+\beta) t] \\ \sigma &= \sqrt{\frac{\sum K_i^2}{2}} V^{-1} \\ \Lambda &= \frac{3(\gamma - \beta - 1)^2}{\gamma^2} H^2 + \frac{\kappa_5^2 \Lambda_5}{2} \left[1 - \frac{(\gamma - \beta - 1)^2}{\gamma^2} \right] \\ G &= \frac{\kappa_5^3 (\gamma - \beta - 1)}{\gamma} \sqrt{\frac{\Lambda_5}{6}} \operatorname{cosech}[\sqrt{\frac{3\Lambda_5}{2}} \kappa_5 (1+\beta) t] \end{aligned} \right\} \quad (37)$$

From this solution we observe that H decreases sharply with time for small t and approaches a constant value $\kappa_5 \sqrt{\frac{\Lambda_5}{6}}$. This assures the positivity of ρ , λ and G . Asymptotically, for large t , the universe expands exponentially with positive acceleration $q \approx -1$ and Λ and G become constant (see figs. 1 and 2) while universe isotropizes due to exponential fall off of the anisotropic scalar (see fig. 3). Also the interdependence of Ω_ρ and β over q has been shown in the 3D graph (fig. 4).

Further, the expressions for density parameter and the age of the universe are

$$\Omega_\rho = \frac{2(\gamma - \beta - 1)(1+q)}{3\gamma^2}$$




Fig.1




Fig.2

Figs. 1 shows variation of G for $\Lambda_5 > 0$ with the choice $\sqrt{\frac{3\Lambda_5}{2}} \kappa_5 = 1$ for $\gamma = 4/3$. Fig. 2 presents the acceleration or deceleration of the universe at different cosmic time for $\Lambda_5 > 0$.




Fig.3




Fig.4

Fig. 3 shows the gradual isotropization of the universe with time. Fig. 4 gives the variation of q with the variation of Ω_ρ and β for $\Lambda_5 \neq 0$.

$$T = \frac{1}{(1+\beta)\sqrt{a}} \tanh^{-1} \left[\frac{2+3\beta-q}{3(1+\beta)} \right]^{1/2} \quad (38)$$

So we can write

$$\beta = \gamma - 1 - \frac{3\Omega_\rho \gamma^2}{2(1+q)} \quad (39)$$

i.e., β can be estimated from the observed values.

Again from the relations (37) we have

$$G = \frac{\kappa_5^2(\gamma - \beta - 1)\sqrt{1+q}}{\gamma\sqrt{3(1+\beta)}} H \quad (40)$$

Also from (30) using (37) we write

$$a = \frac{3}{2} \kappa_5^2 \Lambda_5 = \frac{3H^2(2+3\beta-q)}{1+\beta} \quad (41)$$

Hence from the above two relations we can evaluate the five dimensional quantities κ_5 and Λ_5 . Thus the solution contains no free parameter.

We note that a non-vanishing cosmological constant in the bulk induces a natural time scale in $4D$ as

$$\tau_s = \sqrt{\frac{6}{\kappa_5^2 \Lambda_5}}$$

which in terms of observed quantities has the expression

$$\tau_s = \frac{1}{H} \left(1 - \frac{\gamma^2 \Omega_\rho}{2(\gamma - \beta - 1)(1 + \beta)} \right)^{-1/2} \quad (42)$$

If we consider a model universe with $\Omega_\rho \approx 0.3$ and that of $H \approx 0.7 \times 10^{-10} \text{ yr}^{-1}$, then the characteristic scale is $\approx 5.2 \times 10^{10} \text{ yrs}$ i.e., 52 billion years, which is much greater than the age of the universe (due to data from WMAP mission) [21].

Lastly, from the observational bound for β the deceleration parameter is restricted by $\frac{3\gamma\Omega_\rho}{2} - 1 < q < \frac{3\Omega_\rho\gamma^2}{2(\gamma-.034)} - 1$, which shows that for $\Omega_\rho < \frac{2(\gamma-.034)}{3\gamma^2}$, there is always an accelerating universe while for $\Omega_\rho > \frac{2(\gamma-.034)}{3\gamma^2}$ both acceleration and deceleration is possible (see fig. 4), but for $\Omega_\rho > \frac{2}{3\gamma}$, only we have decelerating universe.

Thus in the present matter dominated era ($\gamma = 1$) to get accelerated universe $\Omega_\rho < 0.644$ which supports the observed value of Ω_ρ . Therefore it is possible to have at present an accelerating universe in this model.

Case II: Λ_5 (i.e., a) = 0

Here the relation (34) should be satisfied and V has the simple solution

$$V = \left[\sqrt{b}(\beta + 1)t \right]^{\frac{1}{\beta+1}}, \quad (\beta + 1 \neq 0) \quad (43)$$

with

$$\xi = \frac{3(\gamma - 1 - \beta)^2}{\gamma^2}$$

(Note that $\beta + 1 \neq 0$; otherwise there is no variation of G)

Here also the integration constant t_0 is taken to be zero for simplicity. The scale factors will have an explicit forms as

$$a_i = a_{i0} V^{\frac{1}{3}} \exp \left[-\frac{K_i}{3\beta\sqrt{b}} V^\beta \right], \quad i = 1, 2, 3$$

The deceleration parameter will have constant value and is related to β by the relation

$$q = 2 + 3\beta$$

The brane tension, $4D$ effective cosmological term Λ and the gravitational coupling G has the time variation as

$$\lambda = \frac{2(\gamma - 1 - \beta)}{\kappa_5^2 \gamma (\beta + 1)t}, \quad \Lambda = \frac{3(\gamma - 1 - \beta)^2}{\gamma^2} H^2, \quad G = \frac{\kappa_5^2 (\gamma - 1 - \beta) H}{\gamma} \quad (44)$$

with

$$H = \frac{1}{3(1 + \beta)t}$$




Fig.5




Fig.6

Figs. 5 and 6 present the variation of q due to the variation of Ω_ρ and γ for '+' and '-' sign in equation (46).

as Hubble parameter.

The anisotropy scalar σ decreases with time as $t^{-\frac{1}{1+\beta}}$ and vanishes at infinity. Now for energy density namely $\rho = \frac{2}{\kappa_5^2 \gamma t}$, we have the following relation among density parameter Ω_ρ , β and γ

$$\frac{G\rho}{3H^2} = \Omega_\rho = \frac{2(\gamma - \beta - 1)(\beta + 1)}{\gamma^2} \quad (45)$$

which gives β as a function of γ and Ω_ρ

$$\beta + 1 = \frac{\gamma}{2} [1 \pm \sqrt{1 - 2\Omega_\rho}] \quad (46)$$

Thus for real β we must have $\Omega_\rho \leq 1/2$. This upper limit of Ω_ρ is independent of any specific values of γ and q . Also eliminating ' β ' between (45) and the expression for q we have

$$\Omega_\rho = \frac{2(3\gamma - q - 1)(1 + q)}{9\gamma^2} \quad (47)$$

The variation of q over Ω_ρ and γ has been shown in figures 5 and 6 for positive and negative signs in equation (46). In the first case there is always deceleration while in the second case q varies between -1 and 2 .

Now the relationship between the density parameters are

$$\frac{\Lambda}{3H^2} = \Omega_\Lambda = \frac{9\gamma^2\Omega_\rho^2}{4(1 + q)^2} \quad (48)$$

Further we note that the above solution can be recovered from the solution for $\Lambda_5 > 0$ if we proceed to the limit $\Lambda_5 \rightarrow 0$. Near the big bang epoch (i.e., for small t) the present solution has similar behaviour with that for positive Λ_5 . Lastly, to have an accelerated universe the density parameter should have further restriction on upper bound (depending on γ), considering '-' sign in equation (46) as

$$\Omega_\rho < \frac{2(3\gamma - 1)}{9\gamma^2} \quad (49)$$

We note that this upper bound for Ω_ρ is always less than 0.5, hence it is possible to have a decelerating phase of the universe when Ω_ρ lies between the bound in equation (49) and 0.5. On the other hand, for ‘+’ sign in equation (46) the universe will accelerate if $\gamma < 2/3$ and then there is a lower bound for Ω_ρ i.e.,

$$\frac{2(3\gamma - 1)}{9\gamma^2} < \Omega_\rho \leq 0.5 \quad (50)$$

To have an end of this section we remark that similar to the previous solution for $\Lambda_5 > 0$, here also there is no free parameter remain, all of them can be estimated from observation (i.e., Ω_ρ and H). In table I and II we have presented the numerical values of different parameters based on various values of the density parameter Ω_ρ for dust model ($\gamma = 1$) with $H = 0.7 \times 10^{-10} \text{ yr}^{-1}$. Table I corresponds to ‘-’ sign in the expression for β in equation (46) while table II for ‘+’ sign. In table I, throughout there is acceleration except for $\Omega_\rho = 0.5$ as expected from the expression for q . In fact, for $\Omega_\rho > 0.44$ only deceleration is possible. We note that here $\Omega_\rho + \Omega_\Lambda \neq 1$, there is a contribution of Ω_{ρ^2} in brane scenario. But this contribution is insignificant in the present epoch. Thus the data in this table are consistent with recent observation as there is deceleration followed by acceleration. On the otherhand, table II has the reverse picture. There is always deceleration as $\gamma = 1$ and Ω_{ρ^2} has significant contribution in the matter content of the universe. Finally, we say that table II represents early phases of the universe when brane scenario has dominant contribution while table I is valid in the present epoch when brane effect is negligible.

TABLE-I

($\Lambda_5 = 0$. Present phase of the evolution, accelerated expansion)

Ω_ρ	β	q	Ω_Λ	$T/10^{10} \text{ yr's}$
0.5	-0.5	0.5	0.25	0.95238
0.4	-0.72361	-0.17082	0.52361	1.72287
0.3	-0.81623	-0.44868	0.66623	2.59120
0.2	-0.88730	-0.66189	0.7873	4.22523
0.1	-0.94721	-0.84164	0.89721	9.02108
0.08	-0.95826	-0.87477	0.91826	11.4078
0.04	-0.97958	-0.93875	0.95958	23.3234

TABLE-II

($\Lambda_5 = 0$. Early phase of the evolution, decelerated expansion)

Ω_ρ	β	q	Ω_Λ	$T/10^{10}$ yr's
0.5	-0.5	0.5	0.25	0.95238
0.4	-0.27639	1.17082	0.07639	0.65808
0.3	-0.18377	1.44868	0.03377	0.58340
0.2	-0.11270	1.66190	0.01270	0.53668
0.1	-0.05279	1.84164	0.00279	0.50273
0.08	-0.04174	1.87477	0.00174	0.49693
0.04	-0.02042	1.93875	0.00042	0.48612

Case III: Λ_5 (i.e., a) < 0

This case is important from the point of view of brane scenario as proposed by Randall and Sundrum [8] because according to them a single brane is embedded in a five dimensional anti-de-Sitter bulk (AdS₅).

For this choice the solution of V can be written as (with $t_0 = 0$)

$$V^{1+\beta} = \sqrt{\frac{b}{|a|}} \sin[(1+\beta)\sqrt{|a|} t] \quad (51)$$

with the expressions for the physical parameters as

$$\left. \begin{aligned} H &= \frac{\theta}{3} = \sqrt{|a|} \cot[(1+\beta)\sqrt{|a|} t] \\ \rho &= \frac{\sqrt{6|\Lambda_5|} (1+\beta)}{\kappa_5 \gamma} \operatorname{cosec}[(1+\beta)\sqrt{|a|} t] \\ q &= 3(1+\beta) \sec^2[(1+\beta)\sqrt{|a|} t] - 1 \\ \lambda &= \frac{(\gamma-\beta-1)\sqrt{6|\Lambda_5|}}{\kappa_5 \gamma} \operatorname{cosec}[(1+\beta)\sqrt{|a|} t] \\ \sigma &= \sqrt{\frac{\sum K_i^2}{2}} V^{-1} \\ \Lambda &= \frac{3(\gamma-\beta-1)^2}{\gamma^2} H^2 + \frac{\kappa_5^2 \Lambda_5}{2} \left[1 - \frac{(\gamma-\beta-1)^2}{\gamma^2} \right] \\ G &= \frac{\kappa_5^3 (\gamma-\beta-1)}{\gamma} \sqrt{\frac{|\Lambda_5|}{6}} \operatorname{cosec}[(1+\beta)\sqrt{|a|} t] \end{aligned} \right\} \quad (52)$$

This solution represents a spatially flat but recollapsing universe with recollapsing time (τ_{rec}) given by

$$\sin[(1+\beta)\sqrt{|a|} \tau_{rec}] = 0$$

Though this solution is completely different from the solution with positive Λ_5 but we can obtain this solution from (37) if we change $\sqrt{\Lambda_5} \rightarrow i\sqrt{|\Lambda_5|}$. As a consequence, $\sqrt{a} \rightarrow i\sqrt{|a|}$ and $\sinh(\sqrt{a} t) \rightarrow i \sin(\sqrt{|a|} t)$. Finally, the age of the universe can be written as

$$\tau = \frac{\tau_{rec}}{\pi} \tan^{-1} \left(\frac{q-2-3\beta}{3(1+\beta)} \right)^{1/2}$$

IV. BIANCHI V MODEL

The Bianchi V space-time is an isotropic generalization of the open ($k = -1$) Robertson-Walker geometry and the metric-ansatz has the expression

$$ds^2 = -dt^2 + a_1^2(t)dx^2 + a_2^2(t)e^{-2x}dy^2 + a_3^2(t)e^{-2x}dz^2 \quad (53)$$

As before the field equations are equation (20) and

$$\frac{1}{V} \frac{d}{dt}(VH_i) - 2a_1^{-2} = \Lambda - \frac{(\gamma-2)}{2} \kappa^2 \rho - \frac{(\gamma-1)}{12} \kappa_5^4 \rho^2 + \frac{U_0}{V^{\frac{4}{3}}}, \quad i = 1, 2, 3 \quad (54)$$

$$2H_1 - H_2 - H_3 = 0 \quad (55)$$

Now compare to the Bianchi I model in the previous section, we have the same relation (28) among the Hubble parameters but the constants are given by

$$K_1 = 0, \quad K_2 = -K_3 \quad (56)$$

due to the field equation (46). The differential equation for V in the present model is of the form (after integrating once)

$$\dot{V}^2 = aV^2 + bV^{-2\beta} + cV^{\frac{2}{3}} - \frac{4}{V} + d_1 \quad (57)$$

Also as before considering the variation of Λ as $\Lambda = \xi H^2$, we have the same differential equation (34) as in Bianchi I model. Comparing (48) with (34) we note that Bianchi V model is possible only for $\beta = \frac{1}{2}$ (with $c = 0 = d_1$) which contradicts the observational bounds on β . The explicit solution is (choosing the integration constants appropriately)

$$t = \begin{cases} \frac{2}{3\sqrt{a}} \cosh^{-1} V^{\frac{3}{2}} \sqrt{\left| \frac{a}{b-4} \right|}, & \text{if } \frac{b-4}{a} < 0 \\ \frac{2}{3\sqrt{a}} \sinh^{-1} V^{\frac{3}{2}} \sqrt{\left| \frac{a}{b-4} \right|}, & \text{if } \frac{b-4}{a} > 0 \\ \frac{2}{3\sqrt{a}} \log V^{3/2}, & \text{if } b = 4 \end{cases} \quad (58)$$

The explicit form for the physical parameters are :

$$\left. \begin{aligned} H &= \frac{\sqrt{a}}{3} \tanh\left(\frac{3}{2}\sqrt{a} t\right) \\ \rho &= D\sqrt{\frac{a}{b-4}} (\cosh\frac{3}{2}\sqrt{a} t)^{-1} \\ q &= -1 - \frac{9}{2} \operatorname{cosech}^2\left(\frac{3}{2}\sqrt{a} t\right) \\ \lambda &= \frac{2}{3}D\left(\gamma - \frac{3}{2}\right)\sqrt{\frac{a}{b-4}} (\cosh\frac{3}{2}\sqrt{a} t)^{-1} \\ \Lambda &= \xi H^2 \\ G &\propto (\cosh\frac{3}{2}\sqrt{a} t)^{-1} \end{aligned} \right\} \quad \text{for } \frac{b-4}{a} < 0 \quad (59)$$

$$\left. \begin{aligned} H &= \frac{\sqrt{a}}{3} \coth\left(\frac{3}{2}\sqrt{a} t\right) \\ \rho &= D \sqrt{\frac{a}{b-4}} (\sinh \frac{3}{2}\sqrt{a} t)^{-1} \\ q &= -1 + \frac{9}{2} \operatorname{sech}^2\left(\frac{3}{2}\sqrt{a} t\right) \\ \lambda &= \frac{2}{3}D\left(\gamma - \frac{3}{2}\right) \sqrt{\frac{a}{b-4}} (\sinh \frac{3}{2}\sqrt{a} t)^{-1} \\ \Lambda &= \xi H^2 \\ G &\propto (\sinh \frac{3}{2}\sqrt{a} t)^{-1} \end{aligned} \right\} \text{ for } \frac{b-4}{a} > 0 \quad (60)$$

$$\left. \begin{aligned} H &= \frac{\sqrt{a}}{3} \\ \rho &= D e^{-\frac{3}{2}\sqrt{a} t} \\ q &= -1 \\ \lambda &= \frac{2}{3}D\left(\gamma - \frac{3}{2}\right) e^{-\frac{3}{2}\sqrt{a} t} \\ \Lambda &= \xi H^2 \\ G &\propto e^{-\frac{3}{2}\sqrt{a} t} \end{aligned} \right\} \text{ for } b = 4 \quad (61)$$

V. DISCUSSION

In this paper, we have studied the consequences of the variation of G and Λ in the form $\frac{\dot{G}}{G} \sim H$ and $\Lambda \sim H^2$ on Bianchi I and V cosmological models based on the brane-world scenario. We have obtained solutions in Bianchi I model for positive, zero and negative value of the cosmological constant Λ_5 on the bulk. From the solution with positive Λ_5 one can obtain solution for negative Λ_5 by analytic continuation ($\sqrt{\Lambda_5} \rightarrow i\sqrt{|\Lambda_5|}$) while solution for $\Lambda_5 = 0$ are the limiting solution of (37) as $\Lambda_5 \rightarrow 0$. However, asymptotically, as $t \rightarrow \infty$ the nature of the solutions in three cases are distinct. For $\Lambda_5 > 0$, the universe expands exponentially with constant Λ as $t \rightarrow \infty$, while for $\Lambda_5 = 0$ the universe expands in a power law fashion (for $\beta \neq -1$) with a constant deceleration and all physical parameters become vanishingly small. But for $\Lambda_5 < 0$, the solution is an expanding-collapsing model of the universe i.e., an oscillatory model with finite time period.

For $\Lambda_5 > 0$, H and G fall off sharply from large positive value for small t to a constant value for large t , while Λ approaches to zero asymptotically. Also for $\Lambda_5 = 0$, the behaviour of these parameters are similar but all of them vanish asymptotically. But we have peculiar behaviour of these parameters for negative Λ_5 as there are points of discontinuities at finite t . We have also presented solutions for Bianchi V model with $\Lambda_5 > 0$ only for different signs (including zero) of the constant $\frac{b-4}{a}$, but they are not realistic as they corresponds to $\beta = 0.5$, which is outside the observational bounds on β . Their mathematical form is very similar to the Bianchi I case, with one correction term, proportional to $V^{4/3}$ added to the parametric time equation. Hence the general physical behaviour in both geometries has the similar qualitative features. Lastly, we have shown the behaviour of the shear scalar




Fig.7

Fig. 7 showing σ against t and β where $\sigma = t^{-\frac{\beta+2}{\beta+1}}$ for standard cosmology Bianchi I model.

($\sigma \propto t^{-\frac{\beta+2}{\beta+1}}$) against t and β in fig.7 for standard cosmology Bianchi I model. In standard model σ falls off as polynomial in ' t ' while in brane scenario it falls exponentially. In fact the figures 3 and 7 show the similarity of the anisotropy scalar in brane scenario and in standard cosmological model.

Finally, we like to emphasize that the whole analysis of this paper is independent of any particular choice of the bulk parameters which is a good characteristic for the brane-world models.

Acknowledgement:

The authors are grateful to the referee for his valuable suggestions. U.D. is thankful to C.S.I.R., Govt. of India for awarding a Senior Research Fellowship.

References:

- [1] V. N. Melnikov, "Gravity as a key Problem of the Millennium", Proc. 2000 NASA/JPL Conference on Fundamental Physics in Microgravity, CD-Version, NASA Document D-21522, 4.1-4.7 (2001) (Solvang, CA, USA), *gr-qc/0007067*.
- [2] P. A. M. Dirac, *Nature* **139** 323 (1937); *Proc. Royal Soc. London* **165** 199 (1938).
- [3] D. W. Sciama, *Mon. Not. R. Astron. Soc.* **113** 34 (1953).
- [4] P. Jordan, Scherkraft and Weltall, Friedrich Vieweg and Sohn, Braunschweig (1955).
- [5] C. Brans and R. Dicke, *Phys. Rev.* **124** 925 (1961).
- [6] S. Perlmutter et al, *Astrophys. J.* **483** 565 (1997).
- [7] Y. Fujii, *Gravitation and Cosmology* **6** 107 (2000).
- [8] M. Gasperini, *Phys. Lett. B* **194** 347 (1987); *Class. Quantum Grav.* **5** 521 (1988); K. Freese, F.C. Adams, J.A. Friemann and E. Mottolla, *Nucl. Phys. B* **287** 797 (1987); P.J.E. Peebles and B. Ratra, *Astrophys. J* **325** L17 (1988); W. Chen and Y.S. Wu, *Phys. Rev. D* **41** 695 (1990).
- [9] S. Perlmutter et al, *Nature* (London) **391** 51 (1998); R.R. Caldwell, R. Dave and P.J. Steinhardt, *Phys. Rev. Lett.* **80** 1582 (1998); L.P. Chimento, A. S. Jakubi and D. Pavon, *Phys. Rev. D* **62** 063508 (2000).
- [10] A. Bonanno and M. Reuter, *Phys. Lett.* **527B** 8 (2002).
- [11] I.L. Shapiro, J. Sola, C. Espana-Bonnet and P. Ruiz-Lapente, *Phys. Lett.* **574B** 149 (2003).
- [12] R. G. Vishwakarma, *Class. Quantum Grav.* **19** 4747 (2002).

- [13] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83** 3370 (1999); *Phys. Rev. Lett.* **83** 4690 (1999).
- [14] K. Maeda and D. Wands, *Phys. Rev. D* **62** 124009 (2000).
- [15] M. Sasaki, T. Shiromizu and K. Maeda, *Phys. Rev. D* **62** 024008 (2000); T. Shiromizu, K. Maeda and M. Sasaki, *Phys. Rev. D* **62** 024012 (2000)
- [16] R. Maartens, *Phys. Rev. D* **62** 084023 (2000).
- [17] R. Maartens, “*Geometry and Dynamics of Brane-World*”, *gr-qc/0101059*.
- [18] C. M. Chen, T. Harko and M. K. Mak, *Phys. Rev. D* **64** 044013 (2001); T. Harko and M. K. Mak, *Class. Quantum Grav.* **20** 407 (2003).
- [19] J. Ponce de Leon, “*Brane Universes with Variable G and $\Lambda_{(4)}$* ”, *Class. Quantum Grav.* **20** 5321 (2003); (*gr-qc/0305041*).
- [20] J. Ponce de Leon, *Mod. Phys. Lett. A* **17** 2425 (2002); *Int. J. Mod. Phys. D* **11** 1355 (2002).
- [21] D. N. Spergel et al, “*First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters*”, *astro-ph/0302209*.